

# ALLEGRA ALLERGY 180-MG TABLETS

## Drug Facts

### Active ingredient (in each tablet)

Fexofenadine HCl 180 mg..... Antihistamine

### Purpose

### Uses

temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:

- runny nose
- sneezing
- itchy, watery eyes
- itching of the nose or throat

### Warnings

**Do not use** if you have ever had an allergic reaction to this product or any of its ingredients.

**Ask a doctor before use if you have** kidney disease. Your doctor should determine if you need a different dose.

### When using this product

- do not take more than directed
- do not take at the same time as aluminum or magnesium antacids
- do not take with fruit juices (see Directions)

**Stop use and ask a doctor if** an allergic reaction to this product occurs. Seek medical help right away.

**If pregnant or breast-feeding**, ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

### Directions

adults and children 12 years of age and over	take one 180 mg tablet with water once a day; do not take more than 1 tablet in 24 hours
children under 12 years of age	do not use
adults 65 years of age and older	ask a doctor
consumers with kidney disease	ask a doctor

### Other information

- safety sealed: do not use if carton is opened or if printed foil inner seal on bottle is torn or missing
- store between 20° and 25°C (68° and 77°F)
- protect from excessive moisture

### Inactive ingredients

colloidal silicon dioxide, croscarmellose sodium, hypromellose, iron oxide blends, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, pregelatinized starch, titanium dioxide

### Questions or comments?

call toll-free 1-800-633-1610 or [www.allegra.com](http://www.allegra.com)

[ALLEGRA DFL.]



# ALLEGRA ALLERGY GELCAPS 24HR

## Drug Facts

### Active ingredient (in each tablet)

Fexofenadine HCl 180 mg ..... Antihistamine

### Purpose

### Uses

temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:

- runny nose
- sneezing
- itchy, watery eyes
- itching of the nose or throat

### Warnings

**Do not use** if you have ever had an allergic reaction to this product or any of its ingredients.

**Ask a doctor before use if you have** kidney disease. Your doctor should determine if you need a different dose.

### When using this product

- do not take more than directed
- do not take at the same time as aluminum or magnesium antacids
- do not take with fruit juices (see Directions)

**Stop use and ask a doctor if** an allergic reaction to this product occurs. Seek medical help right away.

**If pregnant or breast-feeding**, ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

### Directions

adults and children 12 years of age and over	take one 180 mg tablet with water once a day; do not take more than 1 tablet in 24 hours
children under 12 years of age	do not use
adults 65 years of age and older	ask a doctor
consumers with kidney disease	ask a doctor

### Other information

- safety sealed: do not use if carton is opened or if printed foil inner seal on bottle is torn or missing
- store between 20° and 25°C (68° and 77°F)
- protect from excessive moisture

### Inactive ingredients

colloidal silicon dioxide, croscarmellose sodium, hypromellose, iron oxide blends, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, pregelatinized starch, titanium dioxide

### Questions or comments?

call toll-free 1-800-633-1610 or [www.allegra.com](http://www.allegra.com)

[ALLEGRA DFL.]



## ALLEGRA ALLERGY 60-MG TABLETS

### Drug Facts

#### Active ingredient (in each tablet)

Fexofenadine HCl 60 mg ..... Antihistamine

#### Purpose

### Uses

temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:

- runny nose
- sneezing
- itchy, watery eyes
- itching of the nose or throat

### Warnings

**Do not use** if you have ever had an allergic reaction to this product or any of its ingredients.

**Ask a doctor before use if you have** kidney disease. Your doctor should determine if you need a different dose.

#### When using this product

- do not take more than directed
- do not take at the same time as aluminum or magnesium antacids
- do not take with fruit juices (see Directions)

**Stop use and ask a doctor if** an allergic reaction to this product occurs. Seek medical help right away.

**If pregnant or breast-feeding**, ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

### Directions

adults and children 12 years of age and over	take one 60 mg tablet with water every 12 hours; do not take more than 2 tablets in 24 hours
children under 12 years of age	do not use
adults 65 years of age and older	ask a doctor
consumers with kidney disease	ask a doctor

### Other information

- safety sealed: do not use if carton is opened or if printed foil inner seal on bottle is torn or missing
- store between 20° and 25°C (68° and 77°F)
- protect from excessive moisture

### Inactive ingredients

colloidal silicon dioxide, croscarmellose sodium, hypromellose, iron oxide blends, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, pregelatinized starch, titanium dioxide

### Questions or comments?

call toll-free 1-800-633-1610 or [www.allegra.com](http://www.allegra.com)

[ALLEGRA DFL.]



# ALLEGRA-D 24HR ALLERGY & CONGESTION TABLETS

## Drug Facts

### Active ingredients (in each tablet)

Fexofenadine HCl 180 mg .....	Antihistamine
Pseudoephedrine HCl 240 mg .....	Nasal decongestant

### Purpose

### Uses

- temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
  - runny nose
  - itching of the nose or throat
  - sneezing
  - itchy, watery eyes
- temporarily relieves congestion due to the common cold, hay fever or other upper respiratory allergies
- reduces swelling of nasal passages
- temporarily relieves sinus congestion and pressure
- temporarily restores freer breathing through the nose

### Warnings

#### Do not use

- if you have ever had an allergic reaction to this product or any of its ingredients
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product
- if you have difficulty swallowing

#### Ask a doctor before use if you have

- heart disease
- high blood pressure
- trouble urinating due to an enlarged prostate gland
- kidney disease. Your doctor should determine if you need a different dose.
- thyroid disease
- diabetes
- glaucoma

### When using this product

#### do not take more than directed

- do not take at the same time as aluminum or magnesium antacids
- do not take with fruit juices (see Directions)
- the tablet coating may be seen in the stool (this is normal). Continue to take as directed (see Directions).

#### Stop use and ask a doctor if

- an allergic reaction to this product occurs. Seek medical help right away.
- symptoms do not improve within 7 days or are accompanied by a fever
- you get nervous, dizzy, or sleepless

**If pregnant or breast-feeding**, ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

### Directions

- do not divide, crush, chew or dissolve the tablet; swallow tablet whole

adults and children 12 years of age and over	take 1 tablet with a glass of water every 24 hours on an empty stomach; do not take more than 1 tablet in 24 hours
children under 12 years of age	do not use
adults 65 years of age and older	ask a doctor
consumers with kidney disease	ask a doctor

### Other information

- each tablet contains: **sodium 33 mg**
- safety sealed: do not use if carton is opened or if individual blister units are torn or opened
- store between 20° and 25°C (68° and 77°F)

### Inactive ingredients

acetone, black iron oxide, cellulose acetate, colloidal silicon dioxide, copovidone, croscarmellose sodium, FD&C blue #1 aluminum lake, glycerol triacetate, hypromellose, isopropyl alcohol, magnesium stearate, methyl alcohol, methylene chloride, microcrystalline cellulose, polyethylene glycol, propylene glycol, povidone, sodium chloride, talc, titanium dioxide, water

### Questions or comments?

call toll-free 1-800-633-1610 or [www.allegra.com](http://www.allegra.com)



[ALLEGRA DFL.]

## ALLEGRA-D 12HR ALLERGY & CONGESTION TABLETS

### Drug Facts

#### Active ingredients (in each tablet)

Active ingredients (in each tablet)	Purpose
Fexofenadine HCl 60 mg	Antihistamine
Pseudoephedrine HCl 120 mg	Nasal decongestant

#### Uses

- temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
  - runny nose
  - itching of the nose or throat
  - sneezing
  - itchy, watery eyes
- temporarily relieves congestion due to the common cold, hay fever or other upper respiratory allergies
- reduces swelling of nasal passages
- temporarily relieves sinus congestion and pressure
- temporarily restores freer breathing through the nose

#### Warnings

##### Do not use

- if you have ever had an allergic reaction to this product or any of its ingredients
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product
- if you have difficulty swallowing

##### Ask a doctor before use if you have

- heart disease
- high blood pressure
- trouble urinating due to an enlarged prostate gland
- kidney disease. Your doctor should determine if you need a different dose.
- thyroid disease
- diabetes
- glaucoma

#### When using this product

- do not take more than directed**
- do not take at the same time as aluminum or magnesium antacids
- do not take with fruit juices (see Directions)
- the tablet coating may be seen in the stool (this is normal). Continue to take as directed (see Directions).

#### Stop use and ask a doctor if

- an allergic reaction to this product occurs. Seek medical help right away.
- symptoms do not improve within 7 days or are accompanied by a fever
- you get nervous, dizzy, or sleepless

If pregnant or breast-feeding, ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

#### Directions

- do not divide, crush, chew or dissolve the tablet; swallow tablet whole

adults and children 12 years of age and over	take 1 tablet with a glass of water every 12 hours on an empty stomach; do not take more than 2 tablets in 24 hours
children under 12 years of age	do not use
adults 65 years of age and older	ask a doctor
consumers with kidney disease	ask a doctor

#### Other information

- safety sealed: do not use if carton is opened or if individual blister units are torn or opened
- store between 20° and 25°C (68° and 77°F)


#### Inactive ingredients

carnauba wax, colloidal silicon dioxide, croscarmellose sodium, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, pregelatinized starch, stearic acid

#### Questions or comments?

call toll-free 1-800-633-1610 or [www.allegra.com](http://www.allegra.com)

[ALLEGRA DFL.]



## CHILDREN'S ALLEGRA ALLERGY ORAL SUSPENSION

### Drug Facts

Active ingredient (in each 5 mL teaspoonful)	Purpose
Fexofenadine HCl 30 mg .....	Antihistamine

### Uses

temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:

- runny nose
- sneezing
- itchy, watery eyes
- itching of the nose or throat

### Warnings

**Do not use** if you have ever had an allergic reaction to this product or any of its ingredients.

**Ask a doctor before use if you have** kidney disease. Your doctor should determine if you need a different dose.

### When using this product

- do not take more than directed
- do not take at the same time as aluminum or magnesium antacids
- do not take with fruit juices (see Directions)

**Stop use and ask a doctor if** an allergic reaction to this product occurs. Seek medical help right away.

**If pregnant or breast-feeding**, ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

### Directions

- shake well before using
- use only with enclosed dosing cup

adults and children 12 years of age and over	take 2 teaspoonfuls (10 mL) every 12 hours; do not take more than 4 teaspoonfuls (20 mL) in 24 hours
children 2 to under 12 years of age	take 1 teaspoonful (5 mL) every 12 hours; do not take more than 2 teaspoonfuls (10 mL) in 24 hours
children under 2 years of age	ask a doctor
adults 65 years of age and older	ask a doctor
consumers with kidney disease	ask a doctor

Note: teaspoonful=tsp

### Other information

- each 5 mL teaspoon contains: **sodium 18 mg**
- safety sealed: do not use if carton was opened or if printed foil inner seal on bottle is torn or missing
- store between 20° and 25°C (68° and 77°F)

### Inactive ingredients

butylparaben, edetate disodium, flavor, poloxamer 407, propylene glycol, propylparaben, purified water, sodium phosphate dibasic heptahydrate, sodium phosphate monobasic monohydrate, sucrose, titanium dioxide, xanthan gum, xylitol

### Questions or comments?

call toll-free 1-800-633-1610 or [www.allegra.com](http://www.allegra.com)

[ALLEGRA DFL.]



## CHILDREN'S ALLEGRA ALLERGY ORALLY DISINTEGRATING TABLETS

### Drug Facts

#### Active ingredient (in each tablet)

Fexofenadine HCl 30 mg

#### Purpose

Antihistamine

### Uses

temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:

- runny nose
- sneezing
- itchy, watery eyes
- itching of the nose or throat

### Warnings

**Do not use** if you have ever had an allergic reaction to this product or any of its ingredients.

**Ask a doctor before use if you have** kidney disease. Your doctor should determine if you need a different dose.

#### When using this product

- do not take more than directed
- do not take at the same time as aluminum or magnesium antacids
- do not take with fruit juices (see Directions)

**Stop use and ask a doctor if** an allergic reaction to this product occurs. Seek medical help right away.

**If pregnant or breast-feeding**, ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

### Directions

- place 1 tablet on tongue; tablet disintegrates, with or without water

adults and children 12 years of age and over

take 2 tablets every 12 hours on an empty stomach; do not take more than 4 tablets in 24 hours

children 6 to under 12 years of age

take 1 tablet every 12 hours on an empty stomach; do not take more than 2 tablets in 24 hours

children under 6 years of age

do not use

adults 65 years of age and older

ask a doctor

consumers with kidney disease

ask a doctor

### Other information

- each tablet contains: **sodium 5 mg**
- phenylketonurics: contains phenylalanine 5.3 mg per tablet
- safety sealed: do not use if carton is opened or if individual blister units are torn or opened
- store between 20° and 25°C (68° and 77°F)
- use tablet immediately after opening individual blister

### Inactive ingredients

aspartame, citric acid anhydrous, crospovidone, flavors, magnesium stearate, mannitol, methacrylic acid copolymer, microcrystalline cellulose, povidone, sodium bicarbonate, sodium starch glycolate

### Questions or comments?

call toll-free 1-800-633-1610 or [www.allegra.com](http://www.allegra.com)

[ALLEGRA DFL.]

